

Supplementary table 1. Patient characteristics and intraoperative data for the patients with and without significant SAH

	Total (N = 163)	Patients without significant SAH (n = 133)	Patients with significant SAH (n = 30)	P value
Age (years)	72.0 ± 5.1	71.8 ± 5.1	73.2 ± 5.1	0.155
Height (cm)	166.3 ± 6.2	166.5 ± 6.3	165.4 ± 6.0	0.393
Weight (kg)	65.6 ± 8.3	65.3 ± 8.3	66.5 ± 8.7	0.490
BMI (kg/m ²)	23.7 ± 2.4	23.5 ± 2.3	24.5 ± 2.6	0.036
ASA physical status				0.121
I	39 (23.9)	36 (27.1)	3 (10.0)	
II	105 (64.4)	83 (62.4)	22 (73.3)	
III	19 (11.7)	14 (10.5)	5 (16.7)	
Past history				
Cerebrovascular disease	17 (10.4)	13 (9.8)	4 (13.3)	0.565
Myocardial infarction	12 (7.4)	8 (6.0)	4 (13.3)	0.166
Hypertension	83 (50.9)	65 (48.9)	18 (60.0)	0.271
Diabetes mellitus	33 (20.2)	27 (20.3)	6 (20.2)	0.970
Pulmonary disease	24 (14.7)	19 (14.3)	5 (16.7)	0.747
Renal disease	5 (3.1)	4 (3.0)	1 (3.3)	0.926
Liver disease	5 (3.1)	4 (3.0)	1 (3.3)	0.926
Dose of bupivacaine (mg)	12.0 (11.0-12.0)	12.0 (12.0-12.0)	12.0 (12.0-12.0)	0.210
Level of sensory block	T10 (T8-T10)	T10 (T8-T10)	T8 (T6-T10)	0.002

Data are mean ± standard deviation, median (interquartile range), or number (%). SAH, spinal anesthesia-induced hypotension; BMI, body mass index; ASA, American Society of Anesthesiologists.